arXiv:1704.04395v1 [math.FA] 14 Apr 2017

BOUNDED OPERATORS TO ℓ-KÖTHE SPACES

ELIF UYANIK AND MURAT H. YURDAKUL

ABSTRACT. For Fréchet spaces E and F we write $(E,F)\in\mathcal{B}$ if every continuous linear operator from E to F is bounded. Let ℓ be a Banach sequence space with a monotone norm in which the canonical system (e_n) is an unconditional basis. We obtain a necessary and sufficient condition for $(E,F)\in\mathcal{B}$ when $F=\lambda^{\ell}(B)$. We say that a triple (E,F,G) has the bounded factorization property and write $(E,F,G)\in\mathcal{BF}$ if each continuous linear operator $T:E\longrightarrow G$ that factors over F is bounded. We extend some results in [3] to ℓ-Köthe spaces and obtain a sufficient condition for $(E,\lambda^{\ell_1}(A)\hat{\otimes}_{\pi}\lambda^{\ell_2}(B),\lambda^{\ell_3}(C))\in\mathcal{BF}$ when $\lambda^{\ell_1}(A)$ and $\lambda^{\ell_2}(B)$ are nuclear.

Dedicated to the memory of Prof. Dr. Tosun Terzioğlu

1. Introduction

For an infinite matrix $A=(a_n^k)$ with $0\leq a_n^k\leq a_n^{k+1}$ and $\sup\limits_k a_n^k>0$ for every n and k we denote by $\lambda(A)$ the corresponding l_1-Köthe space, that is,

$$\lambda(A)=\left\{x=(x_n):\|x\|_k=\sum_n|x_n|\,a_n^k<\infty,\quad\forall k\in\mathbb{N}\right\}$$

Equipped with the system of seminorms $\{\|.\|_k,k\in\mathbb{N}\}$, $\lambda(A)$ is a Fréchet space.

Following [2], we denote by ℓ a Banach sequence space in which the canonical system (e_n) is an unconditional basis. The norm $\|.\|$ is called monotone if $\|x\|\leq\|y\|$ whenever $|x_n|\leq|y_n|$, $x=(x_n)$, $y=(y_n)\in\ell$, $n\in\mathbb{N}$. Let Λ be the class of such spaces with monotone norm. In particular, $l_p\in\Lambda$ and $c_0\in\Lambda$. It is known that every Banach space with an unconditional basis has a monotone norm which is equivalent

2010 *Mathematics Subject Classification.* 46A03, 46A45, 46A32, 46A04.

Key words and phrases. bounded operators, bounded factorization property, ℓ-Köthe spaces.

This research was partially supported by Turkish Scientific and Technological Research Council.

to its original norm. Indeed, it is enough to put

$$\|x\| = \sup_{|\beta_n|\leq 1} \left| \sum_n {e_n}'(x)\beta_n e_n \right|$$

where $|.|$ denotes the original norm, $({e_n}')$ denote the sequence of coefficient functionals.

Let $\ell \in \Lambda$ and $\|.\|$ be a monotone norm in ℓ. If $A = (a_n^k)$ is a Köthe matrix, the ℓ-Köthe space $\lambda^\ell(A)$ is the space of all sequences of scalars (x_n) such that $(x_n a_n^k) \in \ell$ with the topology generated by the seminorms

$$\|(x_n)\|_k = \left\|(x_n a_n^k)\right\|$$

Let us remind that $\|(e_n)\|_k = a_n^k$. We denote by $\mathcal{L}(E,F)$ the space of all continuous linear operators between Fréchet spaces E and F. For $T \in \mathcal{L}(E,F)$ we consider the following operator seminorms

$$|T|_{p,q} = \sup\left\{ |Tx|_p : |x|_q \leq 1 \right\}, \quad p,q \in \mathbb{N}$$

which may take the value $+\infty$. In particular, for any one dimensional operator $T = u \otimes x$ which sends each $z \in E$ to $u(z)x \in F$, we have

$$|T|_{p,q} = |u|_q^* \, |x|_p$$

where $\|u\|_q^* = \sup\left\{ |u(x)| : \|x\|_q \leq 1 \right\}$.

We recall that $T : E \longrightarrow F$ is continuous if there is a map $N : \mathbb{N} \longrightarrow \mathbb{N}$ such that

$$\|T\|_{k,N(k)} < \infty, \quad \forall k \in \mathbb{N};$$

T is bounded if $\exists N \in \mathbb{N}$ such that

$$\|T\|_{r,N} < \infty, \quad \forall r \in \mathbb{N}.$$

We write $(E,F) \in \mathcal{B}$ if every continuous linear operator from E to F is bounded. For Fréchet spaces E and F, in [4], Vogt proved that $(E,F) \in \mathcal{B}$ if and only if for every sequence $N(k)$, $\exists N \in \mathbb{N}$ such that $\forall r \in \mathbb{N}$ we have $k_0 \in \mathbb{N}$ and $C > 0$ with

$$\|T\|_{r,N} \leq C \max_{1\leq k\leq k_0} \|T\|_{k,N(k)} \tag{1.1}$$

for all $T \in \mathcal{L}(E,F)$.

We say that a triple (E,F,G) has the bounded factorization property and write $(E,F,G) \in \mathcal{BF}$ if each continuous linear operator $T : E \longrightarrow G$ that factors over F is bounded. In [3], the property $\mathcal{BF}$ is characterized not only for triples of Köthe spaces but also for the general case of Fréchet spaces. Our aim here to extend some results in [3] to ℓ-Köthe space case.

2. BOUNDED OPERATORS TO ℓ-KOTHE SPACES

If we follow the steps of Crone and Robinson Theorem [1], we obtain the following.

Lemma 2.1. *$T \in \mathcal{L}(\lambda(A), \lambda^{\ell}(B))$ iff $\forall m$, $\exists k$ such that*

$$\sup_n \frac{\|Te_n\|_m}{\|e_n\|_k} < +\infty$$

Proof. $T \in \mathcal{L}(\lambda(A), \lambda^{\ell}(B))$ iff $\forall m$, $\exists k$ such that

$$\sup_{x\neq 0, x\in\lambda(A)} \frac{\|Tx\|_m}{\|x\|_k} < +\infty$$

For $x = e_n$, we obtain the result.
Conversely, suppose that $\forall m$, $\exists k$ such that

$$\sup_n \frac{\|Te_n\|_m}{\|e_n\|_k} < +\infty$$

Let $x \in \lambda(A)$.

$$\begin{aligned} \|Tx\|_m &= \left\| \sum_n x_n Te_n \right\|_m \leq \sum_n |x_n| \frac{\|Te_n\|_m}{\|e_n\|_k} \|e_n\|_k \\ &\leq \sup_n \frac{\|Te_n\|_m}{\|e_n\|_k} \sum_n |x_n| \, a_n^k \leq \sup_n \frac{\|Te_n\|_m}{\|e_n\|_k} \|x\|_k \end{aligned}$$

So, $T \in \mathcal{L}(\lambda(A), \lambda^{\ell}(B))$. □

Notice that when domain is ℓ-Köthe space, we can not use this argument.

Our first result is the following.

Theorem 2.2. *The following are equivalent:*
i) $(\lambda(A), \lambda^{\ell}(B)) \in \mathcal{B}$
ii) for every sequence $N(k)$, there is $N \in \mathbb{N}$ such that for each $r \in \mathbb{N}$ we have $k_0 \in \mathbb{N}$ and $C > 0$ with

$$\frac{b_v^r}{a_i^N} \leq C \max_{1\leq k\leq k_0} \frac{b_v^k}{a_i^{N(k)}}$$

for all $v \in \mathbb{N}$,$i \in \mathbb{N}$.

Proof. Suppose that $(\lambda(A), \lambda^\ell(B)) \in \mathcal{B}$. Consider $S : \lambda(A) \longrightarrow \lambda^\ell(B)$ with $S = e_i' \otimes e_v$ where $e_i'(x) = x_i$ for all $x \in \lambda(A)$. Since S is the operator of rank one, we note that

$$\|S\|_{k,N(k)} = \|e_i'\|_{N(k)} \|e_v\|_k = \frac{b_v^k}{a_i^{N(k)}}$$

Similarly, $\|A\|_{r,N} = \frac{b_v^r}{a_i^N}$. So, the result follows from (1.1).

For the converse, let $Te_i = \sum_{v=1}^{\infty} u_{vi} e_v$. Since T is continuous, by Lemma 2.1, there is $N(k)$ such that

$$\begin{aligned} \|T\|_{k,N(k)} &= \sup_{i\in\mathbb{N}} \frac{\|Te_i\|_k}{\|e_i\|_{N(k)}} \\ &= \sup_{i\in\mathbb{N}} \sup_{|\beta_v|\leq 1} \left| \sum_{v=1}^{\infty} u_{vi} \beta_v \frac{b_v^k}{a_i^{N(k)}} e_v \right| < \infty \end{aligned}$$

So we find $N \in \mathbb{N}$ such that

$$\begin{aligned} \|T\|_{r,N} &\leq \sup_{i\in\mathbb{N}} \left\{ \sup_{|\beta_v|\leq 1} \left| \sum_{v=1}^{\infty} u_{vi} \beta_v \frac{b_v^r}{a_i^N} e_v \right| \right\} \\ &\leq \sup_{i\in\mathbb{N}} \left\{ \sup_{|\beta_v|\leq 1} \left| \sum_{v=1}^{\infty} u_{vi} \beta_v \left(C \max_{1\leq k\leq k_0} \frac{b_v^k}{a_i^{N(k)}} \right) e_v \right| \right\} \\ &\leq C \sum_{k=1}^{k_0} \sup_{i\in\mathbb{N}} \left\{ \sup_{|\beta_v|\leq 1} \left| \sum_{v=1}^{\infty} u_{vi} \beta_v \frac{b_v^k}{a_i^{N(k)}} e_v \right| \right\} < \infty \end{aligned}$$

Therefore, T is bounded. □

Now, consider the ℓ-Köthe space $\lambda^\ell(A)$ and any Fréchet space E. Then, we obtain the following.

Theorem 2.3. *The following are equivalent:*
i) $(E, \lambda^\ell(A)) \in \mathcal{B}$
ii) for every sequence $N(k)$*, there is* $N \in \mathbb{N}$ *such that for each* $r \in \mathbb{N}$ *we have* $k_0 \in \mathbb{N}$ *and* $C > 0$ *with*

$$a_v^r \|u\|_N^* \leq C \max_{1\leq k\leq k_0} a_v^k \|u\|_{N(k)}^*$$

for all $v \in \mathbb{N}$, $u \in E'$.

Proof. Suppose that $(E, \lambda^\ell(A)) \in \mathcal{B}$. Similar to the proof of Theorem 2.2, consider the operator of rank one $A = y \otimes e_v$ where $y \in E'$. The result follows from (1.1).

For the converse, let $T : E \longrightarrow \lambda^{\ell}(A)$ be continuous linear operator. Let

$$Tx = \sum_{v=1}^{\infty} e_v'(Tx)e_v = (e_v'T(x)) = (u_v(x)), \quad x \in E$$

where $u_v = e_v' \circ T$.

Then, by continuity we find $N(k)$ such that

$$\sup_{\|x\|_{N(k)} \leq 1} \left(\sup_{|\beta_v| \leq 1} \left| \sum_v \beta_v u_v(x) a_v^k e_v \right| \right) = M(k)$$

Let $|u_v(x)| = \theta_v u_v(x)$ where $\theta_v = \pm 1$ and $\beta_v \theta_v = \alpha_v$, note that

$$\begin{aligned}
\|Tx\|_r &= \sup_{|\beta_v| \leq 1} \left| \sum_v \beta_v u_v(x) a_v^r e_v \right| \\
&\leq \sup_{|\beta_v| \leq 1} \left| \sum_v \beta_v \left(\frac{|u_v(x)|}{\|x\|_N} \right) a_v^r e_v \right| \|x\|_N \\
&\leq \sup_{|\beta_v| \leq 1} \left| \sum_v \beta_v \|u_v\|_N^* a_v^r e_v \right| \|x\|_N \\
&\leq \sup_{|\beta_v| \leq 1} \left| \sum_v \beta_v \left(C \max_{1 \leq k \leq k_0} a_v^k \|u_v\|_{N(k)}^* \right) e_v \right| \|x\|_N \\
&\leq C \sum_{k=1}^{k_0} \sup_{|\beta_v| \leq 1} \left| \sum_v \beta_v a_v^k \left(\sup_{\|x\|_{N(k)} \leq 1} |u_v(x)| \right) e_v \right| \|x\|_N \\
&\leq C \sum_{k=1}^{k_0} \sup_{\|x\|_{N(k)} \leq 1} \left(\sup_{|\beta_v| \leq 1} \left| \sum_v \beta_v a_v^k \theta_v u_v(x) e_v \right| \right) \|x\|_N \\
&\leq C \sum_{k=1}^{k_0} \sup_{\|x\|_{N(k)} \leq 1} \left(\sup_{|\alpha_v| \leq 1} \left| \sum_v \alpha_v a_v^k u_v(x) e_v \right| \right) \|x\|_N \\
&\leq \left(C \sum_{k=1}^{k_0} M(k) \right) \|x\|_N
\end{aligned}$$

Hence T is bounded. □

3. BOUNDED FACTORIZATION PROPERTY FOR ℓ-KOTHE SPACES

We need the following theorem [3, Theorem 2.2].

Theorem 3.1. *For Fréchet spaces E, F and G we have $(E, F, G) \in \mathcal{BF}$ if and only if for every sequence $N(k)$ there is $N \in \mathbb{N}$ such that for each $r \in \mathbb{N}$ we have $k_0 = k_0(r) \in \mathbb{N}$ and $C = C(r) > 0$ so that the following inequality*

$$\|T\|_{r,N} \leq C \max_{1\leq k\leq k_0} \left\{ \|R\|_{k,N(k)} \right\} \max_{1\leq k\leq k_0} \left\{ \|S\|_{k,N(k)} \right\}$$

is satisfied for every $R \in L(F, G)$, $S \in L(E, F)$ where $T = RS$.

The next result is obtained by following the lines of [3, Corollary 3.1].

Theorem 3.2. *Let E be a Fréchet space, $\lambda^{\ell}(B), \lambda^{\tilde{\ell}}(C)$ be ℓ-Köthe spaces and $\lambda^{\ell}(B)$ be nuclear. Then $(E, \lambda^{\ell}(B), \lambda^{\tilde{\ell}}(C)) \in \mathcal{BF}$ if and only if for every sequence $N(k)$ there is $N \in \mathbb{N}$ such that for each $r \in \mathbb{N}$ we have $k_0 \in \mathbb{N}$ and $C > 0$ with*

$$c_j^r \|u\|_N^* \leq C \max_{1\leq k\leq k_0} \left\{ \|u\|_{N(k)}^* \, b_i^k \right\} \max_{1\leq k\leq k_0} \left\{ \frac{c_j^k}{b_i^{N(k)}} \right\} \tag{3.1}$$

for all $i \in \mathbb{N}, j \in \mathbb{N}$ and $u \in E'$.

Proof. Let $S = u \otimes e_i$ and $R = e_i' \otimes e_j$ where $u \in E'$. Then $RS : E \longrightarrow G$ is the operator of rank one which sends each $x \in E$ to $u(x)e_j$. If we apply Theorem 3.1 we obtain the result.

For sufficiency, we take $S \in L(E, \lambda^{\ell}(B))$, $R \in L(\lambda^{\ell}(B), \lambda^{\tilde{\ell}}(C))$ and $T = RS$. Since $\lambda^{\ell}(B)$ is nuclear, $\exists S(k)$ such that $S(k) > N(k)$ and

$$\sum_i \frac{b_i^{N(k)}}{b_i^{S(k)}} = \theta(k) < \infty, \quad k \in \mathbb{N}.$$

We can write $Sx = \sum_i u_i(x)e_i$ where $u_i = e_i' \circ S \in E'$ and $Re_i = \sum_j r_{ji}e_j$. Therefore

$$Tx = \sum_i \sum_j u_i(x) r_{ji} e_j$$

For this $S(k)$ we choose $N \in \mathbb{N}$ such that for each $r \in \mathbb{N}$ we obtain $k \in \mathbb{N}$ and $C > 0$ with

$$c_j^r \|u_i\|_N^* \leq C \max_{1\leq k\leq k_0} \left\{ \|u_i\|_{S(k)}^* \, b_i^k \right\} \max_{1\leq k\leq k_0} \left\{ \frac{c_j^k}{b_i^{S(k)}} \right\} \tag{3.2}$$

for all $i \in \mathbb{N}, j \in \mathbb{N}, u_i \in E'$.

Since all types nuclear Köthe spaces determined by one and the same matrix B coincide [2, Corollary 2, p.22], $\lambda(B) = \lambda^{\ell}(B)$ and we have

$$\|S\|_{k,N(k)} = \sup_{\|x\|_{N(k)}\leq 1} \|Sx\|_k = \sup_{\|x\|_{N(k)}\leq 1} \sum_i |u_i(x)|\, b_i^k = \sum_i \|u_i\|^*_{N(k)}\, b_i^k$$

$$\|R\|_{k,N(k)} = \sup_i \frac{\|Re_i\|_k}{\|e_i\|_{N(k)}} = \sup_i \sup_{|\beta_j|\leq 1} \left| \sum_j \frac{r_{ji}\beta_j c_j^k e_j}{b_i^{N(k)}} \right|$$

Therefore, we have

$$\sum_i \|u_i\|^*_{S(k)}\, b_i^k \leq \sum_i \|u_i\|^*_{N(k)}\, b_i^k = \|S\|_{k,N(k)}$$

and

$$\sum_j \frac{|r_{ji}|\, c_j^k}{b_i^{S(k)}} \leq \theta(k) \sup_j \frac{|r_{ji}|\, c_j^k}{b_i^{N(k)}} \leq \theta(k)\, \|R\|_{k,N(k)}$$

(see [2, proof of Corollary 2, p.22]) Hence, by (3.2) we obtain that

$$\begin{aligned}
\|Tx\|_r &\leq \sum_i \sum_j \frac{|u_i(x)|}{\|x\|_N} |r_{ji}|\, c_j^r\, \|x\|_N \\
&\leq \sum_i \sum_j \|u_i\|^*_N\, |r_{ji}|\, c_j^r\, \|x\|_N \\
&\leq \sum_i \sum_j |r_{ji}| \left\{ C \max_{1\leq k\leq k_0} \left\{ \|u_i\|^*_{S(k)}\, b_i^k \right\} \max_{1\leq k\leq k_0} \left\{ \frac{c_j^k}{b_i^{S(k)}} \right\} \right\} \|x\|_N \\
&\leq C \sum_{k=1}^{k_0} \sum_i \|u_i\|^*_{S(k)}\, b_i^k \sum_j \frac{|r_{ji}|\, c_j^k}{b_i^{S(k)}}\, \|x\|_N \\
&\leq C \sum_{k=1}^{k_0} \sum_i \|u_i\|^*_{S(k)}\, b_i^k \theta(k)\, \|R\|_{k,N(k)}\, \|x\|_N \\
&\leq \left(C \sum_{k=1}^{k_0} \theta(k)\, \|S\|_{k,N(k)}\, \|R\|_{k,N(k)} \right) \|x\|_N
\end{aligned}$$

Therefore, T is bounded. □

Recall that projective tensor product of two l_1-Köthe spaces $\lambda(A)$ and $\lambda(B)$ is isomorphic to $\lambda(D)$ where $d_{vz}^k = a_v^k b_z^k$.

Theorem 3.2 enables us to get:

Theorem 3.3. *Suppose* $(E, \lambda^{\ell_1}(A)) \in \mathcal{B}$ *and* $(\lambda^{\ell_2}(B), \lambda^{\ell_3}(C)) \in \mathcal{B}$ *where* E *is a Fréchet space,* $\lambda^{\ell_3}(C)$ *is an* ℓ*-Köthe space,* $\lambda^{\ell_1}(A)$ *and* $\lambda^{\ell_2}(B)$ *are nuclear* ℓ*-Köthe spaces. Then* $(E, \lambda^{\ell_1}(A)\hat{\otimes}_\pi \lambda^{\ell_2}(B), \lambda^{\ell_3}(C)) \in \mathcal{BF}$.

Proof. Given $N(k)$ which is assumed to be non-decreasing. Since $\lambda^{\ell_2}(B)$ is nuclear and $(\lambda^{\ell_2}(B), \lambda^{\ell_3}(C)) \in \mathcal{B}$, we obtain that $\lambda^{\ell_2}(B) = \lambda(B)$ and by Theorem 2.2, $\exists n \in \mathbb{N}$ such that $\forall r \in \mathbb{N}$ we have $k_0 = k_0(r) \in \mathbb{N}$ and $C_1 = C_1(r) > 0$ with

$$\frac{c_j^r}{b_i^n} \leq C_1 \max_{1\leq k\leq k_0} \frac{c_j^k}{b_i^{N(k)}}$$

for all $i \in \mathbb{N}$, $j \in \mathbb{N}$.

We then determine $S(k)$ such that $S(k) = N(n)$ if $1 \leq k \leq n$ and $S(k) > N(k)$ if $n+1 \leq k \leq s_0$. Since $(E, \lambda^{\ell_1}(A)) \in \mathcal{B}$ by Theorem 2.3, for this $S(k)$, we find $m \in \mathbb{N}$ such that $\forall q \in \mathbb{N}$ we have $s_0 = s_0(q)$ and $C_2 = C_2(q)$

$$\begin{aligned} a_v^q \left\|u\right\|_m^* &\leq C_2 \max_{1\leq k\leq s_0} a_v^k \left\|u\right\|_{S(k)}^* \\ &\leq C_2 \max_{n\leq k\leq s_0} a_v^k \left\|u\right\|_{N(k)}^* \end{aligned}$$

Therefore, for this $N(k)$ we have $\tilde{s_0} = s_0(N(k))$ and $\tilde{C_2} = C_2(N(k))$ with

$$\begin{aligned} c_j^r \left\|u\right\|_m^* &= \frac{c_j^r}{b_i^n} b_i^n \left\|u\right\|_m^* \\ &\leq \left\{ C_1 \max_{1\leq k\leq k_0} \frac{c_j^k}{b_i^{N(k)}} \right\} b_i^n \left\|u\right\|_m^* \\ &\leq \left\{ C_1 \max_{1\leq k\leq k_0} \frac{c_j^k}{a_v^{N(k)} b_i^{N(k)}} \right\} a_v^{N(k)} b_i^n \left\|u\right\|_m^* \\ &\leq \left\{ C_1 \max_{1\leq k\leq k_0} \frac{c_j^k}{a_v^{N(k)} b_i^{N(k)}} \right\} \left\{ \tilde{C_2} \max_{n\leq k\leq \tilde{s_0}} a_v^k \left\|u\right\|_{N(k)}^* b_i^n \right\} \\ &\leq \left\{ C_1 \max_{1\leq k\leq k_0} \frac{c_j^k}{a_v^{N(k)} b_i^{N(k)}} \right\} \left\{ \tilde{C_2} \max_{1\leq k\leq \tilde{s_0}} a_v^k b_i^k \left\|u\right\|_{N(k)}^* \right\} \end{aligned}$$

Let $s = \max\{k_0, \tilde{s_0}\}$ and $C = C(r) = C_1 \tilde{C_2}$. We have proved that $\exists m \in \mathbb{N}$ such that $\forall r \in \mathbb{N}$ we have $s \in \mathbb{N}$ and $C > 0$ with

$$c_j^r \left\|u\right\|_m^* \leq C \max_{1\leq k\leq s} \left\{ \frac{c_j^k}{a_v^{N(k)} b_i^{N(k)}} \right\} \max_{1\leq k\leq s} \left\{ \left\|u\right\|_{N(k)}^* a_v^k b_i^k \right\}$$

for all $j \in \mathbb{N}, v \in \mathbb{N}, i \in \mathbb{N}$ and $u \in E'$.

If $\lambda(A)^{\ell_1}$ and $\lambda^{\ell_2}(B)$ are nuclear ℓ-Köthe spaces, then

$$\lambda^{\ell_1}(A) \hat{\otimes}_\pi \lambda^{\ell_2}(B) \cong \lambda(A) \hat{\otimes}_\pi \lambda(B) \cong \lambda(D)$$

is nuclear where $d_{vi}^k = a_v^k b_i^k$ [2, Corollary 2, p.22].

By Theorem 3.2 we obtain that $(E, \lambda^{\ell_1}(A)\hat{\otimes}_\pi\lambda^{\ell_2}(B), \lambda^{\ell_3}(C)) \in \mathcal{BF}$. □

References

[1] L. Crone, W. Robinson, Diagonal maps and diameters in Köthe spaces, *Israel J. Math.*, **20**(1975), 13–22.

[2] M. M. Dragilev, *Bases in Köthe spaces*, Rostov State University, 1983.

[3] T. Terzioğlu, V. Zahariuta, Bounded factorization property for Fréchet spaces, *Math. Nachr.*, **253**(2003), 81–91.

[4] D. Vogt, Frécheträume, zwischen denen jede stetige lineare Abbildung beschränkt ist, *J. Reine Angew. Math.*, **345**(1983), 182–200.

Elif Uyanik, Department of Mathematics, Middle East Technical University, 06800 Ankara, Turkey

E-mail address: euyanik@metu.edu.tr

Murat H. Yurdakul, Department of Mathematics, Middle East Technical University, 06800 Ankara, Turkey

E-mail address: myur@metu.edu.tr